

University of Dundee

Coupling of groundwater, river flow and rainfall in an upland floodplain

Archer, Nicole ; Ó Dochartaigh, Brigid ; MacDonald, Alan; Bonell, Michael; Black, Andrew; Coles, Neil A.

Publication date:
2014

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Archer, N., Ó Dochartaigh, B., MacDonald, A., Bonell, M., Black, A., & Coles, N. A. (2014). *Coupling of groundwater, river flow and rainfall in an upland floodplain*. Abstract from EGU General Assembly 2014, Vienna, Austria.

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Coupling of groundwater, river flow and rainfall in an upland floodplain

Nicole Archer (1), Brighid Ó Dochartaigh (1), Alan MacDonald (1), Mike Bonell (2), Andrew Black (3), and Neil Coles (4)

(1) British Geological Survey, Murchison House, West Mains Road, Edinburgh EH9 3LA, United Kingdom (nicarc@bgs.ac.uk), (2) UNESCO Centre, University of Dundee, Perth Road, Dundee DD1 4HN, United Kingdom, (3) Geography Department, University of Dundee, Perth Road, Dundee DD1 4HN, United Kingdom, (4) Centre for Ecohydrology, University of Western Australia, 32 Stirling Highway, Crawley WA 6009, Australia

Upland floodplains provide an important function in regulating river flows and controlling the coupling of hillslope runoff with rivers. To investigate the responses of floodplain groundwater to river flows and rainfall events, a small floodplain in an upland area of the River Tweed catchment, Scotland, was characterised using geophysics, 3D geological mapping and hydrogeological testing; and monitoring undertaken from September 2011 to February 2013 of: groundwater levels in five pairs of piezometers; river stage and flow at the upstream and downstream limits of the study site; soil moisture on the adjacent hillslope; and meteorological parameters. Periodical groundwater chemistry and residence data were also collected.

The floodplain aquifer is permeable throughout but partially stratified, comprising dominantly alluvial and glaciofluvial sandy gravels between 8 and 15m interspersed with thin, intermittent layers of low permeability silts, clays and peats. Overlying the gravel aquifer is a partial thin cover of low permeability alluvial silts, and it is underlain dominantly by low permeability glaciolacustrine silts and clays. High permeability solifluction deposits mantle much of the adjacent hillslope and provide a rapid connection to the floodplain aquifer.

The unusually wet year of 2012 provides a good example of how a temperate upland floodplain responds to consistently high rainfall. Statistical analysis and graphical interpretation of groundwater level, rainfall, soil moisture and river stage demonstrates that: 1) dominant groundwater flow within the floodplain is in the same direction as the river, from up-valley to down-valley; 2) soil moisture in the hillslope is strongly correlated with local rainfall, but groundwater across much of the floodplain is more strongly influenced by river stage; except 3) groundwater near the edge of floodplain, which responds more slowly to local rainfall and river stage changes ; and 4) subsurface flow from the hillslope to the floodplain occurs during high rainfall events.

A detailed investigation of three flood events, when the river rose above bank level and flooded adjacent fields and groundwater became artesian in parts of the floodplain, suggests that antecedent moisture conditions can partly explain the differences in groundwater response during different flood events, where high intensity or long duration rainfall can cause saturated soil conditions, reducing soil water storage capacity and hence promoting flood conditions. A conceptual model based on field data of groundwater flow after storm events during antecedent unsaturated and saturated soil conditions is presented.